Brain Research, 576 (1992) 358–359
Elsevier

Author Index

Abe, Y., see Kalubi, B., 287
Ammons, W.S., see Standish, A., 12
Aston-Jones, G., see Van Bockstaele, E.J., 59

Baba, A., see Matsuda, T., 263
Baca, J.M., see Mandler, R.N., 337
Bardram, L., see Rehfeld, J.F., 111
Barillot, J.C., see Zheng, Y., 235
Barillot, J.C., see Zheng, Y., 245
Barun, S., see Dalkara, T., 197
Berg-Johnsen, J. and Langmoen, I.A., Temperature sensitivity of thin unmyelinated fibers in rat hippocampal cortex, 319
Boja, J.W., see Patel, A., 173
Bonagamba, L.G.H., see Machado, B.H., 131
Borghese, C.M., see Keller, E.A., 162
Bowers, B.J. and Wehner, J.M., Adrenalectomy and stress modulate $GABA_A$ receptor function in LS and SS mice, 80
Butera, P.C., Willard, D.M. and Raymond, S.A., Effects of PVN lesions on the responsiveness of female rats to estradiol, 304

Cao, F. and Leung, L.S., Effect of atropine and PCPA on the behavioral modulation of paired-pulse response in the hipocampal CA1 region, 339
Carrer, H.F., see Keller, E.A., 162
Carroll, F.I., see Patel, A., 173
Castañeda, E., see Whishaw, I.Q., 1
Chesselet, M.-F., see Soghomonian, J.-J., 68

Daikoku, S., see Hisano, S., 311
Dalkara, T., Erdemli, G., Barum, S. and Onur, R., Glycine is required for NMDA receptor activation: electrophysiological evidence from intact rat hippocampus, 197
Dawson, I.M., see Hodgkiss, J.P., 157
De Luca, C.J., see Kamen, G., 165
Duvauchelle, C.L., Levitin, M., MacConell, L.A., Lee, L.K. and Ettenberg, A., Opposite effects of prefrontal cortex and nucleus accumbens infusions of flupenthixol on stimulant-induced locomotion and brain stimulation reward, 104

Egger, M.D., see Nowakowski, R.S., 152
Enters, E.K., Pascua, J.R., McDowell, K.P., Kapasi, M.Z., Povlishock, J.T. and Robinson, S.E., Blockade of acute hypertensive response does not prevent changes in behavior or in CSF acetylcholine (ACH) content following traumatic brain injury (TBI), 271
Erdemli, G., see Dalkara, T., 197
Ettenberg, A., see Duvauchelle, C.L., 104
Ewing, S.E., Weber, R.J., Zauner, A. and Plunkett, R.J., Recovery in hemiparkinsonian rats following intrastriatal implantation of activated leukocytes, 42

Feldman, J.L., see Greer, J.J., 355
Fiorino, D., see Whishaw, I.Q., 1
Fonnum, F., see Hassel, B., 120

Gao, Y., see Patel, A., 173
Gonzales, C., see Soghomonian, J.-J., 68
Greer, J.J., Smith, J.C. and Feldman, J.L., Glutamate release and presynaptic action of AP4 during inspiratory drive to phrenic motoneurons, 355
Grunwerg, B.S., see Krauthamer, G.M., 277

Hassel, B., Paulsen, R.E., Johnsen, A. and Fonnum, F., Selective inhibition of glial cell metabolism in vivo by fluorocitrate, 120
Hayes, N.L., see Nowakowski, R.S., 152
Hayes, R.L., see Mao, J., 254
Hay-Schmidt, A. and Mikkelsen, J.D., Demonstration of a neuronal projection from the entopeduncular nucleus to the substantia nigra of the rat, 343
Heckroth, J.A., Terminal distribution of mossy fibers traversing the superior cerebellar peduncle in the mouse, 322
Higuchi, T., Honda, K., Takano, S. and Negoro, H., Estrogen fails to reduce tuberoinfundibular dopaminergic neuronal activity and to cause a prolactin surge in lactating, ovariectomized rats, 143
Hilsted, L., see Rehfeld, J.F., 111
Hisano, S., Li, S., Kagotani, Y. and Daikoku, S., Synaptic associations between oxytocin-containing magnocellular neurons and neurons containing corticotropin-releasing factor in the rat magnocellular paraventricular nucleus, 311
Hochberg, R.B., see Toran-Allerand, C.D., 25
Hodgkiss, J.P., Dawson, I.M. and Kelly, J.S., An intracellular study of the action of NAN-190 on neurons in the dorsal raphe nucleus of the rat, 157
Honda, K., see Higuchi, T., 143

Irifune, M., see Kalubi, B., 287

Johnsen, A., see Hassel, B., 120
Johnsen, A.K., see Kirby, R.F., 348
Joseph, J.A., see Yamagami, K., 327

Kagotani, Y., see Hisano, S., 311
Kalubi, B., Takeda, N., Irifune, M., Ogino, S., Abe, Y., Su-Ling, H., Yamano, M., Matsunaga, T. and Tohyama, M., Nasal mucosa sensitization with toluene diisocyanate (TDI) increases preprotachykinin A (PPTA) and preproCGRP mRNAs in guinea pig trigeminal ganglion neurons, 287
Kamei, J., Kawashima, N. and Kasuya, Y., Role of spleen or spleen products in the deficiency in morphine-induced analgesia in diabetic mice, 139
Kamen, G., Greenstein, S.S. and De Luca, C.J., Lateral dominance and motor unit firing behavior, 165
Kapasi, M.Z., see Enters, E.K., 271
Kasuya, Y., see Kamei, J., 139
Kato, H., Wanaka, A. and Tohyama, M., Co-localization of basic fibroblast growth factor-like immunoreactivity and its receptor mRNA in the rat spinal cord and the dorsal root ganglion, 351
Kawashima, N., see Kamei, J., 139
Keller, E.A., Borghese, C.M., Carrer, H.F. and Ramirez, O.A., The learning capacity of high or low performance rats is related to the hippocampus NMDA-receptors, 162
Kelley, R.O., see Rosenberg, G.A., 203
Kelly, J.S., see Hodgkiss, J.P., 157
Kirby, R.F., Thunhorst, R.L. and Johnson, A.K., Effects of a non-peptide angiotensin receptor antagonist on drinking and blood pressure responses to centrally administered angiotensins in the rat, 348
Kornfeld, M., see Rosenberg, G.A., 203
Krauthamer, G.M., Krol, J.G. and Grunwerg, B.S., Effect of superior colliculus lesions on sensory unit responses in the intralaminar thalamus of the rat, 277
Krol, J.G, see Krauthamer, G.M., 277
Künzle, H. and Unger, J.W., Neuropeptide Y-like immunoreactive neurons in the suprachiasmatic-subparaventricular region in the hedgehog-tenrec, 332
Kusumoto, K., Mackay, K.B. and McCulloch, J., The effect of the kappa-opioid receptor agonist CI-977 in a rat model of focal cerebral ischaemia, 147
Kyle, A.L., see Owusu-Yaw, V., 220

Landgraf, R., see Ludwig, M., 231
Langmoen, I.A., see Berg-Johnsen, J., 319
Lasansky, A., Properties of depolarizing bipolar cell responses to central illumination in salamander retinal slices, 181
Lee, L.K., see Duvauchelle, C.L., 104
Lever, J., see Patel, A., 173
Levitin, M., see Duvauchelle, C.L., 104
Lew, R., see Patel, A., 173
Lewin, A.H., see Patel, A., 173
Li, S., see Hisano, S., 311
Liotta, L.A., see Rosenberg, G.A., 203
Lu, J., see Mao, J., 254
Ludwig, M. and Landgraf, R., Does the release of vasopressin within the supraoptic nucleus of the rat brain depend upon changes in osmolality and Ca^{2+}/K^+?, 231

Brain Research, 576 (1992) 358–359
Elsevier

Author Index

Abe, Y., see Kalubi, B., 287
Ammons, W.S., see Standish, A., 12
Aston-Jones, G., see Van Bockstaele, E.J., 59

Baba, A., see Matsuda, T., 263
Baca, J.M., see Mandler, R.N., 337
Bardram, L., see Rehfeld, J.F., 111
Barillot, J.C., see Zheng, Y., 235
Barillot, J.C., see Zheng, Y., 245
Barun, S., see Dalkara, T., 197
Berg-Johnsen, J. and Langmoen, I.A., Temperature sensitivity of thin unmyelinated fibers in rat hippocampal cortex, 319
Boja, J.W., see Patel, A., 173
Bonagamba, L.G.H., see Machado, B.H., 131
Borghese, C.M., see Keller, E.A., 162
Bowers, B.J. and Wehner, J.M., Adrenalectomy and stress modulate $GABA_A$ receptor function in LS and SS mice, 80
Butera, P.C., Willard, D.M. and Raymond, S.A., Effects of PVN lesions on the responsiveness of female rats to estradiol, 304

Cao, F. and Leung, L.S., Effect of atropine and PCPA on the behavioral modulation of paired-pulse response in the hipocampal CA1 region, 339
Carrer, H.F., see Keller, E.A., 162
Carroll, F.I., see Patel, A., 173
Castañeda, E., see Whishaw, I.Q., 1
Chesselet, M.-F., see Soghomonian, J.-J., 68

Daikoku, S., see Hisano, S., 311
Dalkara, T., Erdemli, G., Barum, S. and Onur, R., Glycine is required for NMDA receptor activation: electrophysiological evidence from intact rat hippocampus, 197
Dawson, I.M., see Hodgkiss, J.P., 157
De Luca, C.J., see Kamen, G., 165
Duvauchelle, C.L., Levitin, M., MacConell, L.A., Lee, L.K. and Ettenberg, A., Opposite effects of prefrontal cortex and nucleus accumbens infusions of flupenthixol on stimulant-induced locomotion and brain stimulation reward, 104

Egger, M.D., see Nowakowski, R.S., 152
Enters, E.K., Pascua, J.R., McDowell, K.P., Kapasi, M.Z., Povlishock, J.T. and Robinson, S.E., Blockade of acute hypertensive response does not prevent changes in behavior or in CSF acetylcholine (ACH) content following traumatic brain injury (TBI), 271
Erdemli, G., see Dalkara, T., 197
Ettenberg, A., see Duvauchelle, C.L., 104
Ewing, S.E., Weber, R.J., Zauner, A. and Plunkett, R.J., Recovery in hemiparkinsonian rats following intrastriatal implantation of activated leukocytes, 42

Feldman, J.L., see Greer, J.J., 355
Fiorino, D., see Whishaw, I.Q., 1
Fonnum, F., see Hassel, B., 120

Gao, Y., see Patel, A., 173
Gonzales, C., see Soghomonian, J.-J., 68
Greer, J.J., Smith, J.C. and Feldman, J.L., Glutamate release and presynaptic action of AP4 during inspiratory drive to phrenic motoneurons, 355
Grunwerg, B.S., see Krauthamer, G.M., 277

Hassel, B., Paulsen, R.E., Johnsen, A. and Fonnum, F., Selective inhibition of glial cell metabolism in vivo by fluorocitrate, 120
Hayes, N.L., see Nowakowski, R.S., 152
Hayes, R.L., see Mao, J., 254
Hay-Schmidt, A. and Mikkelsen, J.D., Demonstration of a neuronal projection from the entopeduncular nucleus to the substantia nigra of the rat, 343
Heckroth, J.A., Terminal distribution of mossy fibers traversing the superior cerebellar peduncle in the mouse, 322
Higuchi, T., Honda, K., Takano, S. and Negoro, H., Estrogen fails to reduce tuberoinfundibular dopaminergic neuronal activity and to cause a prolactin surge in lactating, ovariectomized rats, 143
Hilsted, L., see Rehfeld, J.F., 111
Hisano, S., Li, S., Kagotani, Y. and Daikoku, S., Synaptic associations between oxytocin-containing magnocellular neurons and neurons containing corticotropin-releasing factor in the rat magnocellular paraventricular nucleus, 311
Hochberg, R.B., see Toran-Allerand, C.D., 25
Hodgkiss, J.P., Dawson, I.M. and Kelly, J.S., An intracellular study of the action of NAN-190 on neurons in the dorsal raphe nucleus of the rat, 157
Honda, K., see Higuchi, T., 143

Irifune, M., see Kalubi, B., 287

Johnsen, A., see Hassel, B., 120
Johnsen, A.K., see Kirby, R.F., 348
Joseph, J.A., see Yamagami, K., 327

Kagotani, Y., see Hisano, S., 311
Kalubi, B., Takeda, N., Irifune, M., Ogino, S., Abe, Y., Su-Ling, H., Yamano, M., Matsunaga, T. and Tohyama, M., Nasal mucosa sensitization with toluene diisocyanate (TDI) increases preprotachykinin A (PPTA) and preproCGRP mRNAs in guinea pig trigeminal ganglion neurons, 287
Kamei, J., Kawashima, N. and Kasuya, Y., Role of spleen or spleen products in the deficiency in morphine-induced analgesia in diabetic mice, 139
Kamen, G., Greenstein, S.S. and De Luca, C.J., Lateral dominance and motor unit firing behavior, 165
Kapasi, M.Z., see Enters, E.K., 271
Kasuya, Y., see Kamei, J., 139
Kato, H., Wanaka, A. and Tohyama, M., Co-localization of basic fibroblast growth factor-like immunoreactivity and its receptor mRNA in the rat spinal cord and the dorsal root ganglion, 351
Kawashima, N., see Kamei, J., 139
Keller, E.A., Borghese, C.M., Carrer, H.F. and Ramirez, O.A., The learning capacity of high or low performance rats is related to the hippocampus NMDA-receptors, 162
Kelley, R.O., see Rosenberg, G.A., 203
Kelly, J.S., see Hodgkiss, J.P., 157
Kirby, R.F., Thunhorst, R.L. and Johnson, A.K., Effects of a non-peptide angiotensin receptor antagonist on drinking and blood pressure responses to centrally administered angiotensins in the rat, 348
Kornfeld, M., see Rosenberg, G.A., 203
Krauthamer, G.M., Krol, J.G. and Grunwerg, B.S., Effect of superior colliculus lesions on sensory unit responses in the intralaminar thalamus of the rat, 277
Krol, J.G, see Krauthamer, G.M., 277
Künzle, H. and Unger, J.W., Neuropeptide Y-like immunoreactive neurons in the suprachiasmatic-subparaventricular region in the hedgehog-tenrec, 332
Kusumoto, K., Mackay, K.B. and McCulloch, J., The effect of the kappa-opioid receptor agonist CI-977 in a rat model of focal cerebral ischaemia, 147
Kyle, A.L., see Owusu-Yaw, V., 220

Landgraf, R., see Ludwig, M., 231
Langmoen, I.A., see Berg-Johnsen, J., 319
Lasansky, A., Properties of depolarizing bipolar cell responses to central illumination in salamander retinal slices, 181
Lee, L.K., see Duvauchelle, C.L., 104
Lever, J., see Patel, A., 173
Levitin, M., see Duvauchelle, C.L., 104
Lew, R., see Patel, A., 173
Lewin, A.H., see Patel, A., 173
Li, S., see Hisano, S., 311
Liotta, L.A., see Rosenberg, G.A., 203
Lu, J., see Mao, J., 254
Ludwig, M. and Landgraf, R., Does the release of vasopressin within the supraoptic nucleus of the rat brain depend upon changes in osmolality and Ca^{2+}/K^+?, 231

Machado, B.H. and Bonagamba, L.G.H., Microinjection of L-glutamate into the nucleus tractus solitarii increases arterial pressure in conscious rats, 131
Mackay, K.B., see Kusumoto, K., 147
MacLusky, N.J., see Toran-Allerand, C.D., 25
Malagon, M., Vallarino, M., Tonon, M.C. and Vaudry, H., Localization and charaterization of diazepam-binding inhibitor (DBI)-like peptides in the brain and pituitary of the trout (*Salmo gairdneri*), 208
Mandler, R.N. and Baca, J.M., Muscle lactate dehydrogenase activity is decreased in murine motor neuron disease, 337
Mao, J., Price, D.D., Mayer, D.J., Lu, J. and Hayes, R.L., Intrathecal MK-801 and local nerve anesthesia synergistically reduce nociceptive behaviors in rats with experimental peripheral mononeuropathy, 254
Matsuda, T., Shimizu, I., Murata, Y. and Baba, A., Glucose and oxygen deprivation induces a Ca^{2+}-mediated decrease in (Na^+ + K^+)-ATPase activity in rat brain slices, 263
Matsunaga, T., see Kalubi, B., 287
Mayer, D.J., see Mao, J., 254
Mikkelsen, J.D., see Hay-Schmidt, A, 343
Miranda, R.C., see Toran-Allerand, C.D., 25
Mittleman, G., see Whishaw, I.Q., 1
Mogensen, N.W., see Rehfeld, J.F., 111
Monstein, H., see Rehfeld, J.F., 111
Murata, Y., see Matsuda, T., 263
Muteki, T., see Sugiyama, K., 97

Negoro, H., see Higuchi,T., 143
Nowakowski, R.S., Hayes, N.L. and Egger, M.D., Competitive interactions during dendritic growth: a simple stochastic growth algorithm, 152
Nováková, J., see Říčný, J., 215
Nowakowski, R.S., Hayes, N.L. and Egger, M.D., Competitive interactions during dendritic growth: a simple stochastic growth algorithm, 152

Ogino, S., see Kalubi, B., 287
Ogita, K., see Pingping, Z., 168
Onur, R., see Dalkara, T., 197
Owusu-Yaw, V., Kyle, A.L. and Stell, W.K., Effects of lesions of the optic nerve, optic tectum and nervus terminalis on rod precursor proliferation in the goldfish retina, 220

Pascua, J.R., see Enters, E.K., 271
Patel, A., Boja, J.W., Lever, J., Lew, R., Simantov, R., Carroll, F.I., Lewin, A.H., Philip, A., Gao, Y. and Kuhar, M.J., A cocaine analog and a GBR analog label the same protein in rat striatal membranes, 173
Paulsen, R.E., see Hassel, B., 120
Philip, A., see Patel, A., 173
Pinault, D., Ectopic axonal firing in an epileptic cortical focus is not triggered by thalamocortical volleys during the interictal stage, 175
Pingping, Z., Ogita, K. and Yoneda, Y., Presence the binding of a variety of ligands related to ionotropic excitatory amino acid receptors in rat retina, 168
Plunkett, R.J., see Ewing, S.E., 42
Povlishock, J.T., see Enters, E.K., 271
Price, D.D., see Mao, J., 254

Ramirez, O.A., see Keller, E.A., 162
Raymond, S.A., see Butera, P.C., 304
Rehfeld, J.F., Mogensen, N.W., Bardram, L., Hilsted, L. and Monstein, H., Expression, but failing maturation of procholecystokinin in cerebellum, 111
Říčný, J., Tuček, S and Nováková, J., Acetylcarnitiine, carnitine and glucose diminish the effect of muscarinic antagonist quinuclidinyl benzilate on striatal acetylcholine content 215
Robinson, S.E., see Enters, E.K., 271
Rosenberg, G.A., Kornfeld, M., Estrada, E., Kelley, R.O., Liotta, L.A. and Stetler-Stevenson, W.G., TIMP-2-reduces proteolytic opening of blood–brain barrier by type IV collagenase, 203
Roth, G.S., see Yamagami, K., 327

Salomon, Y., see Zohar, M., 49
Schino, G. and Troisi, A., Opiate receptor blockade in juvenile macaques: effect on affiliative interactions with their mothers and group companions, 125
Schreihofer, A.M., see Sved, A.F., 297
Shimizu, I., see Matsuda, T., 263
Shimoji, K., see Sugiyama, K., 97
Simantov, R., see Patel, A., 173
Smith, J.C., see Greer, J.J., 355
Soghomonian, J.-J., Gonzales, C. and Chesselet, M.-F., Messenger RNAs encoding glutamate-decarboxylases are differentially affected by nigrostriatal lesions in subpopulations of striatal neurons, 68
Standish, A., Vizzard, M.A and Ammons, W.S., Tonic descending modulation of spinal neurons with renal input, 12
Stell, W.K., see Owusu-Yaw, V., 220
Stetler-Stevenson, W.G., see Rosenberg, G.A., 203
Sugiyama, K., Muteki, T. and Shimoji, K., Halothane-induced hyperpolarization and depression of postsynaptic potentials of guinea pig thalamic neurons in vitro, 97
Su-Ling, H., see Kalubi, B., 287
Supko, D.E., Uretsky, N.J. and Wallace, L.J., AMPA/kainic acid glutamate receptor antagonism in the zona incerta dorsal to the subthalamic nucleus inhibits amphetamine-induced stereotypy but not locomotor activity, 89
Sved, A.F., Tsukamoto, K. and Schreihofer, A.M., Stimulation of α_2-adrenergic receptors in nucleus tractus solitarius is required for the baroreceptor reflex, 297

Takeda, N., see Kalubi, B., 287
Takano, S., see Higuchi, T., 143
Thunhorst, R.L., see Kirby, R.F., 348
Tohyama, M., see Kalubi, B., 287
Tohyama, M., see Kato, H., 351
Tonon, M.C., see Malagon, M., 208
Toran-Allerand, C.D., Miranda, R.C., Hochberg, R.B. and MacLusky, N.J., Cellular variations in estrogen receptor mRNA translation in the developing brain: evidence from combined [^{125}I]-estrogen autoradiography and non-isotopic in situ hybridization histochemistry, 25
Troisi, A., see Schino, G., 125
Tsukamoto, K., see Sved, A.F., 297
Tuček, S., see Říčný, J., 215

Unger, J.W., see Künzle, H., 332
Uretsky, N.J., see Supko, D.E., 89

Vallarino, M., see Malagon, M., 208
Van Bockstaele, E.J. and Aston-Jones, G., Distinct populations of neurons in the ventromedial periaqueductal gray project to the rostral ventral medulla and abducens nucleus, 59
Vaudry, H., see Malagon, M., 208
Vizzard, M.A., see Standish, A., 12

Wallace, L.J., see Supko, D.E., 89
Wanaka, A., see Kato, H., 351
Weber, R.J., see Ewing, S.E., 42
Wehner, J.M., see Bowers, B.J., 80
Whishaw, I.Q., Fiorino, D., Mittleman, G. and Castañeda, E., Do forebrain structures complete for behavioral expression? Evidence from amphetamine-induced behavior, microdialysis, and caudate-accumbens lesions in medial frontal cortex damaged rats, 1
Willard, D.M., see Butera, P.C., 304

Yamagami, K., Joseph, J.A. and Roth, G.S., Decrement of muscarinic receptor-stimulated low-K_MGTPase in striatum and hippocampus from the aged rat, 327
Yamano, M., see Kalubi, B., 287
Yoneda, Y., see Pingping, Z., 168

Zauner, A., see Ewing, S.E., 42
Zheng, Y., Barillot, J.C. and Bianchi, A.L., Intracellular electrophysiological and morphological study of the medullary inspiratory neurons of the decerebrate rat, 235
Zheng, Y., Barillot, J.C. and Bianchi, A.L., Medullary expiratory neurons in the decerebrate rat: an intracellular study, 245
Zohar, M. and Salomon, Y., Melanocortins stimulate proliferation and induce morphological changes in cultured rat astrocytes by distinct transducing mechanisms, 49